



**TAGEM**  
AR-GE & İNOVASYON

DİYARBAKIR ZİRAİ MÜCADELE  
ARAŞTIRMA ENSTİTÜSÜ MÜDÜRLÜĞÜ

## DR. ADİL TONGA

Ziraat Yüksek Mühendisi

### EĞİTİM

- Doktora** 2016-2021 Dicle Üniversitesi  
Fen Bilimleri Enstitüsü
- Yüksek Lisans** 2009-2014 Dicle Üniversitesi  
Fen Bilimleri Enstitüsü
- Lisans** 2004-2008 Dicle Üniversitesi  
Ziraat Fakültesi


**YABANCI DİL** İngilizce YÖKDİL 95.00 / YDS 82.50  
Almanca A1 Kurs Sertifikası/2022

### İŞ TECRÜBESİ


- 2013-Halen, Araştırmacı, Diyarbakır Ziraî Mücadele Araştırma Enstitüsü- DİYARBAKIR
- 2016-2017 (6 ay), Araştırmacı, Max Planck Kimyasal Ekoloji Enstitüsü, Almanya, Jena Almanya
- 2009-2013 Mühendis, Tarım ve Orman İl Müdürlüğü - ŞIRNAK

### HAKKIMDA

### İLETİŞİM

 Silvan Karayolu 7. Km. PK.115 21110  
Sur/DİYARBAKIR / TÜRKİYE

 [adil.tonga@tarimorman.gov.tr](mailto:adil.tonga@tarimorman.gov.tr)

 0 412 326 1143 - 138



▪ **YAYINLAR**

▪ **Uluslararası Dergilerde Yayınlanmış Makaleler**

1. **Tonğa, A.**, Şeker, K., Çakmak, S., Temiz, M. G., & Bayram, A. (2022). Cotton treatment with methyl jasmonate at different growth stages reduces the population of sucking insect pests, and marginally increases their associated predators. *Entomologia Experimentalis et Applicata*. **170**: 207-221. <https://doi.org/10.1111/EEA.13137>
2. **Tonğa, A.**, & Bayram, A. (2021). Natural parasitism of maize stemborers, *Sesamia* spp. (Lepidoptera: Noctuidae) eggs by *Trichogramma evanescens* (Hymenoptera: Trichogrammatidae) in Southeastern Turkey. *International Journal of Agriculture Environment and Food Sciences*, 5(2), 197-202. <https://doi.org/10.31015/jaefs.2021.2.9>
3. **Tonğa, A.**, Ataş, M., & Bayram, A. (2021). Maize (*Zea mays* L.) as a new host plant for the naked grass-mealybug, *Heterococcus nudus* (Hemiptera: Pseudococcidae). *Entomological News*, 129(4), 386-394. <https://doi.org/10.3157/021.129.0406>
4. **Tonğa, A.**, Çakmak, S., Şeker, K., Temiz, M. G., & Bayram, A. (2020). cis-Jasmone treatments affect multiple sucking insect pests and associated predators in cotton. *Entomologia Generalis*, 40(1): 49-61. <https://doi.org/10.1127/entomologia/2019/0771>
5. Mohammadi, S., Maroufpoor, N., **Tonğa, A.**, Bayram, A., & Maroufpoor, M. (2020). Comparative demography and population projection of *Ephestia kuehniella* (Lepidoptera: Pyralidae) and *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Journal of Entomological Society of Iran*, 40(2), 167-181. <https://doi.org/10.22117/JESI.2020.342453.1372>
6. Mutlu, Ç., Karaca, V., **Tonğa, A.**, Erol, Ş., and Mamay, M. (2019). Infestation and damage caused by Wheat stem sawflies (Hymenoptera: Cephidae) to some wheat cultivars in Southeast Anatolia Region, Turkey. *Journal of the Kansas Entomological Society*, 92(1), 359-375. <https://doi.org/10.2317/0022-8567-92.1.359>
7. Bayram A., & **Tonğa, A.** (2018). Methyl Jasmonate affects population densities of phytophagous and entomophagous insects in wheat. *Applied Ecology and Environmental Research*. 16(1): 181-198. [http://doi.org/10.15666/aeer/1601\\_181198](http://doi.org/10.15666/aeer/1601_181198)
8. Bayram A., & **Tonğa, A.** (2018). cis-Jasmone treatments affect pests and beneficial insects of wheat (*Triticum aestivum* L.): the influence of doses and plant growth stages. *Crop Protection*. 105:70-79 <https://doi.org/10.1016/j.cropro.2017.11.011>
9. Bayram A., & **Tonğa, A.** (2016). First report of *Chilo partellus* in Turkey, a new invasive maize pest for Europe. *Journal of Applied Entomology*. 140: 236–240. <http://doi.org/10.1111/jen.12232>